

(12) **United States Patent**
Midoux et al.

(10) Patent No.: **US 6,372,499 B1**
(45) Date of Patent: **Apr. 16, 2002**

(54) **POLYMERIC COMPLEXES FOR THE TRANSFECTION OF NUCLEIC ACIDS, WITH RESIDUES CAUSING THE DESTABILISATION OF CELL MEMBRANES**

(75) Inventors: **Patrick Midoux, Orleans; Michel Monsigny, Saint-Cyr-en Val, both of (FR)**

(73) Assignee: **I.D.M. Immuno-Designed Molecules (FR)**

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/297,519**

(22) PCT Filed: **Nov. 10, 1997**

(86) PCT No.: **PCT/FR97/02022**

§ 371 Date: **May 3, 1999**

§ 102(e) Date: **May 3, 1999**

(87) PCT Pub. No.: **WO98/22610**

PCT Pub. Date: **May 28, 1998**

(30) Foreign Application Priority Data

Nov. 15, 1996 (FR) 96 13990

(51) Int. Cl.⁷ **C12N 15/88**

(52) U.S. Cl. **435/455; 435/69.1; 435/320.1; 435/325; 424/486**

(58) Field of Search **424/486; 435/320.1; 435/455, 325, 69.1; 514/44**

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,166,320 A 11/1992 Wu et al. 530/395
5,733,762 A * 3/1998 Midoux et al. 435/172.3
5,846,530 A * 12/1998 Soon-Shiong et al. 424/93.7
5,977,084 A * 11/1999 Szoka, Jr. et al. 514/44
6,030,941 A * 2/2000 Summerton et al. 514/2

FOREIGN PATENT DOCUMENTS

EP 0350246 1/1990
EP 0387775 9/1990
EP 0388758 9/1990
FR 2107756 5/1972
FR 2719316 11/1995
WO 9211037 7/1992
WO 9213570 8/1992
WO 95/30020 * 11/1995

OTHER PUBLICATIONS

Anderson, Human gene therapy, Nature, vol. 392, pp. 25-30, 1998.*

Verma et al., Gene therapy—promises, problems and prospects, Nature, vol. 389, pp. 239-242, 1997.*

Branch, A good antisense molecule is hard to find, TIBS, Feb. 23, 1998, pp. 45-50.*

Erbacher et al., Glycosylated Polylysine/DNA Complexes: Gene Transfer Efficiency in Relation with the Size and the Sugar Substitution Level of Glycosylated Polylysines and with the Plasmid Size, Bioconjugate Chem., 1995, vol. 6, pp. 401-410.*

Erbacher et al., Gene Transfer by DNA/Glycosylated Polylysine Complexes into Human Blood Macrophages, Human Gene Therapy, vol. 7, Apr. 10, 1996, pp. 721-729.*

Kollen et al., Gluconoylated and Glycosylated Polylysines As Vector for Gene Transfer on Cystic Fibrosis Airways Epithelial Cells, Human Gene Therapy, vol. 7, Aug. 20, 1996, pp. 1577-1586.*

Mezo et al., "Carrier . . . Poly (L-Lysine)", Biopolymers, vol. 33, No. 6, 1993, pp. 873-885, XP002034603.

Wang et al., "Polyhistidine . . . Liposomes", Biochemistry, vol. 23, No. 19, 1984, pp. 4409-4416, XP002016041.

Midoux et al., "Specific . . . Cells", Nucleic Acids Research, vol. 21, No. 4, Feb. 25, 1993, pp. 871-878, XP000371764. Biochemistry, 1984, 23, 4409-4416, XP002016041, Wang et al.

Biopolymers, vol. 33, 873-885, MEZO et al, XP002034603, 1993.

* cited by examiner

Primary Examiner—Dave T. Nguyen

(74) Attorney, Agent, or Firm—Bierman, Muserlian and Lucas

(57) **ABSTRACT**


The complex has at least one negatively charged nucleic acid bonded to at least one positively charged polymeric conjugate

The conjugate containing a polylysine formed from monomers having free NH_3^+ groups, and having at least 10% of the free NH_3^+ groups substituted by residues which can be protonated in a weakly acid medium causing destabilization of cell membranes.

Optionally, some of the free NH_3^+ groups can be substituted by a molecule with a recognition signal by a cell membrane receptor.

The free NH_3^+ groups of the said polylysine make up at least 30% of the monomers of the polymeric conjugate.

The residue that causes the destabilization of cell membrane in weak acid of quinolines of the formula:



where R_1 is hydrogen, R_2 is $\text{-(CH}_2\text{)}_n\text{CO}_2\text{-H}$, X is hydrogen or chlorine and n is an integer from 1 to 10.

The signal is a simple oside or a disaccharide or peptide.

15 Claims, 9 Drawing Sheets